



Effects of Fentanyl Versus Remifentanyl Pretreatment on the QTc Interval in Patients Undergoing Rapid Sequence Intubation: A Randomized Clinical Trial

Seyedpouya Paknezhad¹, Elaheh Serati², Robab Mehdizadeh Esfanjani³, Maryam Soleimanpour⁴ and Hassan Soleimanpour^{5,*}

¹Emergency and Trauma Care Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

²Students Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

³Neurosciences Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁴Social Determinants of Health Research Center, Tabriz University of Medical sciences, Tabriz, Iran

⁵Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

*Corresponding author: Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. Email: h_mofid1357@yahoo.com

Received 2022 August 27; Revised 2022 November 05; Accepted 2022 November 16.

Abstract

Background: One of the complications of laryngoscopy is cardiac dysrhythmia due to an increased QT interval.

Objectives: This study aimed to compare the effect of fentanyl and remifentanyl on QTc interval in patients undergoing intubation by the rapid method in 2020.

Methods: In this study, 50 patients without a history of heart disease and without a history of drug use who were candidates for rapid intubation in the emergency department were selected and divided into two groups. Before the injection of intubation drugs, a baseline ECG was obtained from the patient and then randomly assigned to one group of patients with 2 µg/kg fentanyl and the other group with 1 µg/kg remifentanyl. Other intubation drugs, including lidocaine, etomidate, and succinylcholine, were fixed in both groups. Immediately after drug injection and immediately after laryngoscopy, ECG was prepared from patients, and the QT interval was calculated.

Results: In group comparison, changes in QT values during the studied time were statistically significant ($P = 0.004$). However, in the comparison between groups, there was no statistically significant difference between the two groups receiving fentanyl and remifentanyl ($P = 0.235$). The results showed that the rate of QT interval changes during three different measurements increased in both groups of patients receiving fentanyl and remifentanyl.

Conclusions: In the present study, in comparing the QT interval, there was a statistically significant difference between the two groups receiving fentanyl and remifentanyl at any of the studied times. Therefore, fentanyl and remifentanyl during endotracheal intubation did not have a statistically and clinically significant effect on the QT distance of patients' ECG, and the administration of the above two drugs during intubation was safe.

Keywords: QT Interval, Rapid Sequence Intubation, Laryngoscopy, Fentanyl, Remifentanyl

1. Background

Laryngoscopy and endotracheal intubation can increase heart rate and blood pressure, irregular heartbeat, catecholamines, myocardial oxygen demand, myocardial ischemia, and increased intracerebral pressure in susceptible individuals (1).

These changes are well tolerated in patients without cardiac problems, but in patients with cardiovascular problems and high blood pressure, especially uncontrolled, they can cause many complications (including ischemia) and cardiac dysrhythmias (2,3).

The rapid sequence intubation (RSI) method is the cornerstone of modern airway management and includes the preparation of equipment and drugs, supplemental oxygenation, induction, muscle paralysis with a muscle relaxant, endotracheal placement, and postintubation supportive therapies (4). RSI is the standard method of airway control in emergency patients (5). In this method, medication is used to reduce the complications of glottal stimulation during laryngoscopy. One of these complications is the possibility of cardiac dysrhythmia. Increasing the length of the modified QT interval, which indicates the potential for ventricular dysrhythmias, may be associated with de-

veloping dangerous ventricular dysrhythmias, torsade du point at the top (6). Various mechanisms have been mentioned to prolong this interval. One of the above mechanisms is the increase of sympathetic tone, which during laryngoscopy also increases the probability of prolongation of QT interval with increasing sympathetic stimulation (7). Drugs such as fentanyl or remifentanyl can be used to improve this problem. These drugs can reduce sympathetic tone, and this reduction has a protective role against dysrhythmias caused by QT elongation (8). Both fentanyl and remifentanyl are widely used in RSI, but a few studies have compared the effects of these agents on the QT interval.

These studies are all on patients who are candidates for intubation in the operating room, and no study has been performed on emergency patients.

2. Objectives

Due to the lack of studies on emergency patients in this domain and the small number of studies in this field in general, a study was designed to compare the effect of fentanyl and remifentanyl on the QTc interval in patients undergoing RSI.

3. Methods

This is a double-blinded randomized clinical trial performed from 2019 April 21 to 2019 October 23 in the emergency medicine department of Imam Reza General Hospital Tabriz, Iran. The authors confirm that this intervention's ongoing and related trials are registered. Randomization was done using a table of randomization numbers. The sample size was 25 samples in each group.

3.1. Study Population

For this purpose, 50 patients who were candidates for RSI in the emergency department were selected and divided into two groups.

The inclusion criteria were patients older than 18 who needed rapid sequence intubation for airway management. Patients with a history of cardiovascular disease, diabetes mellitus, chronic hypertension, intracranial injury, drug poisoning, and chronic medication use were excluded.

3.2. Study Design

Before starting the RSI process, a baseline ECG was obtained from the patient, and then one group of patients randomly received 2 μ g/kg fentanyl, and the other group

received 1 μ g/kg remifentanyl. Other intubation drugs, including lidocaine, etomidate, and succinylcholine, were injected into both groups. Immediately after drug injection and immediately after laryngoscopy, ECG was obtained, and the QT interval was calculated manually based on ECG traces. Corrected QT (QTc) was also calculated in all patients using the Bazett formula: $QTc = QT / \sqrt{RR}$. The ECG device was the same.

In the first group, 25 patients requiring rapid intubation received 1 mg/kg lidocaine and two mcg/kg fentanyl 5 minutes before laryngoscopy. Three minutes later, 0.2 mg/kg of etomidate and 1.5 mg/kg of succinylcholine were injected, and laryngoscopy and intubation were performed 90 seconds later. All patients underwent a 12-lead ECG one minute after the lidocaine and fentanyl injection before starting the drug injection.

In the second group, 25 patients requiring rapid intubation received 1 mg/kg lidocaine and one mcg/kg remifentanyl 5 minutes before laryngoscopy. Three minutes later, 0.2 mg/kg of etomidate and 1.5 mg/kg of succinylcholine were injected, and laryngoscopy and intubation were performed 90 seconds later.

In both groups, the mean QTc in the three ECGs, the shortest QT interval, the longest QT interval, and the QT interval dispersion in the obtained ECGs were calculated. Then the results of each group before and after drug injection and the results of the two groups were compared.

The nurse who administered the drugs and the investigator who analyzed the ECG traces were blind to the group of enrolled patients.

Figure 1 shows a flow chart of the measurements of these parameters.

3.3. Analysis

Data were analyzed using SPSS 22.0.0 (IBM® Microsoft L.T.D., Chicago) software. Kolmogorov-Smirnov test and skewness and kurtosis indexes were used to test the normality of data distribution. The results were reported in frequency (percent) and mean $\pm$ standard. Other inferential statistical methods such as independent samples *t*-test, Paired samples *t*-test, and repeated measures analysis of variance (Repeated Measurements) were used to analyze the data. $P < 0.05$ was considered a statistically significant level.

3.4. Ethics

This study was approved by the regional ethics committee with No. IR.TBZMED.REC.1397.921 and registered in IRCT with IRCT registration number: IRCT20090923002496N12




Figure 1. Flow chart of the study

4. Results

In this study, 50 patients were referred to the emergency department. The mean age of patients in the Fentanyl group was 60.08 ± 15.81 years, with a minimum age of 31 years and a maximum age of 90 years. The mean age of patients receiving remifentanyl was 61.16 ± 18.68 years, with a minimum age of 24 years and a maximum age of 87 years. There was no statistically significant difference in age between the two groups ($P = 0.826$).

Nineteen (76%) patients in the fentanyl group were male, and six (24%) were female. Eighteen (72%) of the patients receiving remifentanyl were male, and seven (28%) were female. There was no significant difference in terms

of the sex distribution between the two groups of patients ($P = 0.747$).

The means and standard deviations of the QT interval measured 5 minutes before laryngoscopy, 1 minute after injection, and immediately after laryngoscopy are grouped in [Table 1](#). There was no statistically significant difference in any measurements between the two groups receiving fentanyl and remifentanyl ($P = 0.219$, $P = 0.181$, and $P = 0.368$, respectively) ([Table 1](#)).

In the within-group comparison, changes in QT values over time (in three stages of measurement: 5 minutes before laryngoscopy, one minute after drug injection, and immediately after laryngoscopy) were statistically signifi-

Table 1. Between Groups and Within Groups Comparison of QT Values in Different Measurements

	Group		P.v	P.v (Within Group)
	Fentanyl	Remifentanyl		
QT calculation intervals				0.004
5 minutes before laryngoscopy	9.04 ± 1.1	8.6 ± 1.38	0.219	
1 minute after Laryngoscopy	9.2 ± 1.19	8.72 ± 1.31	0.181	
Actually, after laryngoscopy	9.24 ± 1.16	8.92 ± 1.32	0.368	
P.v (between group)		0.235		

cant ($P = 0.004$). Nevertheless, in the between-group comparison, there was no statistically significant difference between the two groups receiving fentanyl and remifentanyl ($P = 0.235$). Table 2 shows changes in QT values.

The rate of change in QT interval during three different measurements increased in both groups of patients receiving fentanyl and remifentanyl. However, this trend did not show a statistically significant difference in pairwise comparison between QT measurements (Table 2).

The trend of changes in QT values by groups receiving fentanyl and remifentanyl in three measurements is shown in Figure 2.

QT values are given separately in groups of fentanyl and remifentanyl in three measurements in Figure 3.

5. Discussion

In the present study, no significant difference was observed in sex distribution and mean age between the two groups of patients. In the comparison of QT interval, no statistically significant difference was observed between the two groups receiving fentanyl and remifentanyl in any of the studied times (5 minutes before laryngoscopy, one minute after drug injection, and immediately after laryngoscopy). In group comparison, changes in QT values during the studied time were statistically significant ($P = 0.004$). But in the comparison between groups, there was no statistically significant difference between the two groups receiving fentanyl and remifentanyl ($P = 0.235$). The results showed that the rate of QT interval changes during three different measurements increased in both groups of patients receiving fentanyl and remifentanyl. However, this trend did not show a statistically significant difference in pairwise comparison between QT measurements. Therefore, fentanyl and remifentanyl during endotracheal intubation did not have a statistically and clinically significant effect on the QT interval of patients' ECG, and the administration of the above two drugs during intubation is safe.

In the study of TD Kweon et al., The results showed a better effect of remifentanyl 1 mg/kg in reducing the QTc inter-

val during intubation (9), and in the study of T. Cafiero et al. decreased risk of dysrhythmia was observed, the amount of decreased QT interval dispersion, which is better related to a decreased risk of dysrhythmias, was higher in remifentanyl group (10). The above results are not in line with the present study's findings.

O'Hare et al. Reported that remifentanyl 0.5 mg/kg was ineffective in controlling increased heart rate and arterial pressure after intubation, but doses of 1.0 and 1.25 mg/kg effectively controlled the response. Therefore, 1.25 mg/kg remifentanyl was associated with a decrease in systolic arterial pressure of less than 90 mm Hg in seven patients (11).

Pouraghaei et al. Reported that no significant differences were observed between fentanyl, alfentanil, and sufentanil in cardiopulmonary responses to intubation; there is not any difference between these three opioids in the hemodynamic state of patients, and all are safe (12), which is consistent with the findings of the present study.


In a similar study by Yang comparing remifentanyl with fentanyl, the prevalence of systolic blood pressure was significantly higher in the fentanyl group than in the remifentanyl group by increasing more than 30% of baseline values during the study period, but the prevalence of systolic blood pressure decreased. < 30% baseline values were significantly lower in the fentanyl group than in the remifentanyl group (13).

In a study comparing the effects of remifentanyl and fentanyl on changes in blood pressure and arrhythmia in patients undergoing coronary artery bypass grafting, they reported that remifentanyl and fentanyl and routine methods were able to inhibit cardiovascular stimulation induced by laryngoscopy and endotracheal intubation. They undergo coronary artery surgery. But remifentanyl was more effective than the other two drugs up to three minutes after intubation. (14). The present study's findings also showed that remifentanyl and fentanyl could inhibit cardiovascular stimulation caused by laryngoscopy and intubation.

In another study, Min et al. Examined the effects of remifentanyl compared to lidocaine on mitigating hemo-

Table 2. Pair Comparison of QT Values in Different Measurements

Group	QT Calculation Intervals		Mean Difference	Pv	Confidence Interval
Fentanyl	Actually, after laryngoscopy	5 minutes before laryngoscopy	0.2	0.171	-0.06 - 0.46
		1 minute after laryngoscopy	0.04	0.982	-0.06 - 0.14
	1 minute after laryngoscopy	5 minutes before laryngoscopy	0.16	0.31	-0.08 - 0.40
Remifentanyl	Actually, after laryngoscopy	5 minutes before laryngoscopy	0.32	0.880	-0.03 - 0.67
		1 minute after laryngoscopy	0.2	0.304	-0.13 - 0.53
	1 minute after laryngoscopy	5 minutes before laryngoscopy	0.12	0.52	-0.05 - 0.29

**Figure 2.** The trend of QT distance changes by group (QT1: 5 minutes before laryngoscopy. QT2: 1 minute after injection. QT3: Immediately after laryngoscopy)

dynamic responses to laryngoscopy and endotracheal intubation and concluded that remifentanyl was more effective than lidocaine in reducing the hemodynamic response to laryngoscopy and endotracheal intubation (15).

In another study, Ozcan et al. compared the effects of the two drugs. In the alfentanil group, mean arterial blood pressure and heart rate increased in the first minute and during skin incision in comparison with the remifentanyl group. While the use of remifentanyl compared to alfentanil makes the removal of the endotracheal tube faster, and it concluded that both drugs could control the cardiovascular response resulting from laryngoscopy and endotracheal intubation (16).

In a study, Ko et al. compared the effects of fentanyl and remifentanyl on cardiovascular changes following laryngoscopy and endotracheal intubation and concluded that both drugs were effective, but remifentanyl was more effective than fentanyl in attenuating the hemodynamic response (17).

5.1. Conclusions

In the present study, in comparing the QT interval, no statistically significant difference was observed between the two groups receiving fentanyl and remifentanyl at any of the studied times (5 minutes before laryngoscopy, one minute after drug injection, immediately after laryngoscopy). Therefore, fentanyl and remifentanyl during endotracheal intubation did not have a statistically and clinically significant effect on the QT interval of patients' ECG, and the administration of the above two drugs during intubation is safe.

Acknowledgments

We are grateful to all who participated in this study. Our deep appreciation goes to data collectors, supervisors, and administrative staff of the Emergency Department of Imam Reza Hospital, Tabriz, Iran. This article was written




Figure 3. QT interval values by a group of patients receiving fentanyl and Remifentanyl (QT1: 5 minutes before laryngoscopy. QT2: 1 minute after injection. QT3: Immediately after laryngoscopy)

based on a dataset of Elaheh Serati's Medical Specialty degree thesis entitled, "Effects of fentanyl versus Remifentanyl pretreatment on the QTc interval in patients undergoing rapid sequence intubation." This study was registered at Tabriz University of Medical Sciences (Code No: 60886). We would like to appreciate the cooperation of the Clinical Research Development Unit, Imam Reza General Hospital, Tabriz, Iran, in conducting this research.

Footnotes

Authors' Contribution: All authors have read and approved the manuscript. SPP, ES, and HS performed the data collection, literature review, and drafting of the manuscript. MS and RME undertook the major parts of the study design and performed the statistical analysis.

Clinical Trial Registration Code: IRCT20090923002496N12 (04/05/2019).

Conflict of Interests: The authors declare no conflict of interest.

Data Reproducibility: The datasets generated during and analyzed during the current study are not publicly available due to restrictions of the ethics committee of Tabriz University of Medical Sciences but are available from the corresponding author upon reasonable request.

Ethical Approval: This study was approved by the regional ethics committee with No. IR.TBZMED.REC.1397.921.

Funding/Support: It was not declared by the authors.

Informed Consent: Written informed consent was obtained from each patient's relatives.

References

- Henderson J. Adverse effect of anesthesia. In: Miller RD, editor. *Miller's Anesthesia*. 7th ed. New York: Churchill living stone Elsevier Company; 2010.
- Nussmeier N, Husser M, Sarwar M, Grigore A. Anesthesia for cardiac surgical procedures. In: Miller RD, editor. *Miller's Anesthesia*. 2. 7th ed. New York: Churchill living stone Elsevier Company; 2010. 1955 p.
- Stoeling R, Dierdorf S. Anesthesia and coexisting disease. In: Miller RD, editor. *Miller's Anesthesia*. 5th ed. New York: Churchill living stone Elsevier Company; 2008. 87 p.
- Vachon CA, Warner DO, Bacon DR. Succinylcholine and the open globe. Tracing the teaching. *Anesthesiology*. 2003;**99**(1):220-3. [PubMed: 12826863]. <https://doi.org/10.1097/0000542-200307000-00033>.
- Walls RM, Hockberger R, Gausche-Hill M. *Rosens emergency medicine concepts and clinical practice*. 9th ed. Philadelphia: Elsevier; 2018.
- Staikou C, Stamelos M, Stavroulakis E. Impact of anaesthetic drugs and adjuvants on ECG markers of torsadogenicity. *Br J Anaesth*. 2014;**112**(2):217-30. [PubMed: 24305646]. <https://doi.org/10.1093/bja/aet412>.
- Cinca J, Evangelista A, Montoyo J, Barutell C, Figueras J, Valle V, et al. Electrophysiologic effects of unilateral right and left stellate ganglion block on the human heart. *Am Heart J*. 1985;**109**(1):46-54. [PubMed: 3966332]. [https://doi.org/10.1016/0002-8703\(85\)90414-4](https://doi.org/10.1016/0002-8703(85)90414-4).
- Korpinen R, Saarnivaara L, Siren K. QT interval of the ECG, heart rate and arterial pressure during anaesthetic induction: comparative effects of alfentanil and esmolol. *Acta Anaesthesiol Scand*. 1995;**39**(6):809-13. [PubMed: 7484039]. <https://doi.org/10.1111/j.1399-6576.1995.tb04175.x>.

9. Kweon TD, Nam SB, Chang CH, Kim MS, Lee JS, Shin CS, et al. The effect of bolus administration of remifentanyl on QTc interval during induction of sevoflurane anaesthesia. *Anaesthesia*. 2008;**63**(4):347-51. [PubMed: [18336483](#)]. <https://doi.org/10.1111/j.1365-2044.2007.05372.x>.
10. Cafiero T, Di Minno RM, Di Iorio C. QT interval and QT dispersion during the induction of anesthesia and tracheal intubation: a comparison of remifentanyl and fentanyl. *Minerva Anesthesiol*. 2011;**77**(2):160-5. [PubMed: [21150850](#)].
11. O'Hare RA, Mirakhur RK, Reid JE, Breslin DS, Hayes A. Recovery from propofol anaesthesia supplemented with remifentanyl. *Br J Anaesth*. 2001;**86**(3):361-5. [PubMed: [11573525](#)]. <https://doi.org/10.1093/bja/86.3.361>.
12. Pouraghaei M, Moharamzadeh P, Soleimanpour H, Rahmani F, Safari S, Mahmoodpoor A, et al. Comparison between the effects of alfentanil, fentanyl and sufentanil on hemodynamic indices during rapid sequence intubation in the emergency department. *Anesth Pain Med*. 2014;**4**(1). e14618. [PubMed: [24660155](#)]. [PubMed Central: [PMC3961020](#)]. <https://doi.org/10.5812/aapm.14618>.
13. Yang QY, Xue FS, Liao X, Liu HP, Luo MP, Xu YC, et al. Comparison of bolus remifentanyl versus bolus fentanyl for blunting cardiovascular intubation responses in children: a randomized, double-blind study. *Chin Med J (Engl)*. 2009;**122**(1):44-50. [PubMed: [19187616](#)].
14. Nasiri E, Soliamani A, Mohammadpour RA, Donyavi R, Jafari H. [Comparison between the effects of remifentanyl and fentanyl on blood pressure and cardiac dysrhythmia during tracheal intubation in CABG]. *J Mazandaran Univ Med Sci*. 2010;**20**(76):24-31. Persian.
15. Min JH, Chai HS, Kim YH, Chae YK, Choi SS, Lee A, et al. Attenuation of hemodynamic responses to laryngoscopy and tracheal intubation during rapid sequence induction: remifentanyl vs. lidocaine with esmolol. *Minerva Anesthesiol*. 2010;**76**(3):188-92. [PubMed: [20203546](#)].
16. Maguire AM, Kumar N, Parker JL, Rowbotham DJ, Thompson JP. Comparison of effects of remifentanyl and alfentanil on cardiovascular response to tracheal intubation in hypertensive patients. *Br J Anaesth*. 2001;**86**(1):90-3. [PubMed: [11575417](#)]. <https://doi.org/10.1093/bja/86.1.90>.
17. Ko BJ, Oh JN, Lee JH, Choi SR, Lee SC, Chung CJ. Comparison of effects of fentanyl and remifentanyl on hemodynamic response to endotracheal intubation and myoclonus in elderly patients with etomidate induction. *Korean J Anesthesiol*. 2013;**64**(1):12-8. [PubMed: [23372880](#)]. [PubMed Central: [PMC3558642](#)]. <https://doi.org/10.4097/kjae.2013.64.1.12>.